

Product Datasheet

Histone H3 [Trimethyl Lys27] Antibody - BSA Free NBP2-59206

Unit Size: 50 ug

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-59206

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-59206



NBP2-59206

Histone H3 [Trimethyl Lys27] Antibody - BSA Free


Product Information	
Unit Size	50 ug
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide and 0.05% ProClin 300
Isotype	IgG
Purity	Peptide affinity purified
Buffer	PBS
Target Molecular Weight	15 kDa

Product Description	
Description	Novus Biologicals Rabbit Histone H3 [Trimethyl Lys27] Antibody - BSA Free (NBP2-59206) is a polyclonal antibody validated for use in WB, ELISA, ICC/IF and ChIP. Anti-Histone H3 Antibody: Cited in 2 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	126961
Gene Symbol	H3C14
Species	Human, Mouse, Rat, A. thaliana, Drosophila, Parasite, Zebrafish
Reactivity Notes	Schistosoma
Immunogen	The exact sequence of the immunogen to this Histone H3 [Trimethyl Lys27] antibody is proprietary.


Product Application Details	
Applications	Western Blot, Dot Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Chromatin Immunoprecipitation (ChIP), Chromatin Immunoprecipitation Sequencing, Knockdown Validated
Recommended Dilutions	Western Blot 1:1000, ELISA 1:200, Immunocytochemistry/ Immunofluorescence 1:500, Dot Blot 1:20000, Chromatin Immunoprecipitation (ChIP) 1 ug/IP, Chromatin Immunoprecipitation Sequencing, Knockdown Validated
Application Notes	Use in KD reported in scientific publication PMID: 32427586

Images


Western Blot: Histone H3 [Trimethyl Lys27] Antibody [NBP2-59206] - Western blot was performed on whole cell (40 ug, lane 1) and histone extracts (15 ug, lane 2) from HeLa cells, and on 1 ug of recombinant histone H2A, H2B, H3 and H4 (lane 3, 4, 5 and 6, respectively) using the antibody against H3K27me3. The antibody was diluted 1:1,000 in TBS-Tween containing 5% skimmed milk. Observed molecular weight is ~15 kDa.




Immunocytochemistry/Immunofluorescence: Histone H3 [Trimethyl Lys27] Antibody [NBP2-59206] - HeLa cells were stained with the antibody against H3K27me3 and with DAPI. Cells were fixed with 4% formaldehyde for 10' and blocked with PBS/TX-100 containing 5% normal goat serum and 1% BSA. The cells were immunofluorescently labeled with the H3K27me3 antibody (left) diluted 1:500 in blocking solution followed by an anti-rabbit antibody conjugated to Alexa Fluor 488. The middle panel shows staining of the nuclei with DAPI. A merge of the two stainings is shown on the right.




Chromatin Immunoprecipitation: Histone H3 [Trimethyl Lys27] Antibody [NBP2-59206] - ChIP assays were performed using human K562 cells, the antibody against H3K27me3 and optimized PCR primer pairs for qPCR. ChIP was performed with a ChIP-seq kit, using sheared chromatin from 100,000 cells. The indicated amounts of antibody were used per ChIP experiment. IgG (1 ug/IP) was used as a negative IP control. Quantitative PCR was performed with primers specific for the promoter of the active genes GAPDH and EIF4A2, used as negative controls, and TSH2B and MYT1, used as positive controls. The figure shows the recovery, expressed as a % of input (the relative amount of immunoprecipitated DNA compared to input DNA after qPCR analysis).




Western Blot: Histone H3 [Trimethyl Lys27] Antibody [NBP2-59206] - Western blot was performed on whole cell (25 ug, lane 1) and histone extracts (15 ug, lane 2) from HeLa cells, and on 1 ug of recombinant histone H2A, H2B, H3 and H4 (lane 3, 4, 5 and 6, respectively) using the antibody against H3K27me3 diluted 1:500 in TBS-Tween containing 5% skimmed milk. Observed molecular weight is ~15 kDa.




Chromatin Immunoprecipitation: Histone H3 [Trimethyl Lys27] Antibody [NBP2-59206] - ChIP assays were performed using human K562 cells, the antibody against H3K27me3 and optimized PCR primer pairs for qPCR. ChIP was performed with a ChIP-seq kit, using sheared chromatin from 1 million cells. The indicated amounts of antibody were used per ChIP experiment. IgG (1 ug/IP) was used as a negative IP control. Quantitative PCR was performed with primers specific for the promoter of the active genes GAPDH and EIF4A2, used as negative controls, and TSH2B and MYT1, used as positive controls. The figure shows the recovery, expressed as a % of input (the relative amount of immunoprecipitated DNA compared to input DNA after qPCR analysis).



ELISA: Histone H3 [Trimethyl Lys27] Antibody [NBP2-59206] - To determine the titer of the antibody, an ELISA was performed using a serial dilution of the antibody against H3K27me3. The antigen used was a peptide containing the histone modification of interest. By plotting the absorbance against the antibody dilution, the titer of the antibody was estimated to be >1:1,000,000.



Dot Blot: Histone H3 [Trimethyl Lys27] Antibody [NBP2-59206] - To test the cross reactivity of the antibody against H3K27me3, a Dot Blot analysis was performed with peptides containing other modifications or unmodified sequences of histone H3 and H4. One hundred to 0.2 pmol of the respective peptides were spotted on a membrane. The antibody was used at a dilution of 1:20,000. Figure shows a high specificity of the antibody for the modification of interest.



Publications

Yang Y, Chen C, Zuo Q Et al. NARF is a hypoxia-induced coactivator for OCT4-mediated breast cancer stem cell specification Sci Adv 2022-12-09 [PMID: 36490339] (Chemotaxis, Human)

Details:

Citation using the DyLight 405 version of this antibody.

Lu H, Xie Y, Tran L et al. Chemotherapy-induced S100A10 recruits KDM6A to facilitate OCT4-mediated breast cancer stemness J Clin Invest. 2020-05-19 [PMID: 32427586] (KD, WB, Human)



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-59206

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-59206

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

