

Product Datasheet

Nrf2 [p Ser40] Antibody (SU0334) NBP2-67465

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 4

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-67465

Updated 2/24/2026 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-67465



NBP2-67465


Nrf2 [p Ser40] Antibody (SU0334)

Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	SU0334
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	TBS (pH7.4), 0.05% BSA, 40% Glycerol
Product Description	
Description	Novus Biologicals Rabbit Nrf2 [p Ser40] Antibody (SU0334) (NBP2-67465) is a recombinant monoclonal antibody validated for use in IHC, WB and Flow. Anti-Nrf2 Antibody: Cited in 3 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	4780
Gene Symbol	NFE2L2
Species	Human, Mouse, Rat
Immunogen	Synthetic phospho-peptide corresponding to residues surrounding Ser40 of Human Nrf2 aa 20-69 / 605. (SwissProt: Q16236 Human)
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Flow Cytometry, Immunohistochemistry
Recommended Dilutions	Western Blot 1:500-1:2,000, Flow Cytometry 1:50-1:100, Immunohistochemistry, Immunohistochemistry-Paraffin 1:50-1:200




Images


Western Blot: Nrf2 [p Ser40] Antibody (SU0334) [NBP2-67465] -
Western blot analysis of Nrf2 on different lysates. Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. The primary antibody (1/500) was used in 5% BSA at room temperature for 2 hours. Goat Anti-Rabb




Flow Cytometry: Nrf2 [p Ser40] Antibody (SU0334) [NBP2-67465] -
Analysis of RIP3 was done on SW620 cells. The cells were fixed, permeabilized and stained with the primary antibody (1/50) (red). After incubation of the primary antibody at room temperature for an hour, the cells were stained with a Alexa Fluor 488-conjugated Goat anti-Rabbit IgG Secondary antibody at 1/1000 dilution for 30 minutes. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).



Immunohistochemistry-Paraffin: Nrf2 [p Ser40] Antibody (SU0334) [NBP2-67465] -Rat kidney tissue using anti-RIP3 antibody. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 8.0-8.4) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (1/50) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



Western Blot: Nrf2 [p Ser40] Antibody (SU0334) [NBP2-67465] -
Analysis of RIP3 on HT-29 cell lysates with Rabbit anti-RIP3 antibody at 1/1,000 dilution. Lysates/proteins at 10 ug/Lane. Predicted band size: 57 kDa Observed band size: 50 kDa Exposure time: 2 minutes; 12% SDS-PAGE gel. Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody at 1/1,000 dilution was used in 5% NFDM/TBST at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1:100,000 dilution was used for 1 hour at room temperature.



Publications

Schwertheim S, Alhardan M, Manka PP et al. Higher pNRF2, SOCS3, IRF3, and RIG1 Tissue Protein Expression in NASH Patients versus NAFL Patients: pNRF2 Expression Is Concomitantly Associated with Elevated Fasting Glucose Levels *Journal of Personalized Medicine* 2023-07-18 [PMID: 37511764]

Yang J, Choi M, Shin H et al. *Phyllostachys nigra* Variety Henosis, a Domestic Bamboo Species, Protects PC12 Cells from Oxidative Stress-mediated Cell Injury through Nrf2 Activation *Biotechnology and Bioprocess Engineering* 2022-06-21

Park JW, Oh JH, Hwang D et al. 3,4,5-Trihydroxycinnamic acid exerts anti-inflammatory effects on TNF alpha/IFN gamma stimulated HaCaT cells *Molecular medicine reports* 2021-07-01 [PMID: 33982762] (WB, Human)

De Blasio A, Di Fiore R, Pratelli G et al. A loop involving NRF2, miR-29b-1-5p and AKT, regulates cell fate of MDA-MB-231 triple-negative breast cancer cells *J. Cell. Physiol.* 2019-07-10 [PMID: 31313842]





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-67465

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-67465

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

