

Product Datasheet

Collagen VI alpha 1 Antibody (SD83-03) NBP2-67825

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-67825

Updated 2/24/2026 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-67825



NBP2-67825

Collagen VI alpha 1 Antibody (SD83-03)


Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	SD83-03
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	TBS (pH7.4), 0.05% BSA, 40% Glycerol

Product Description	
Description	Novus Biologicals Rabbit Collagen VI alpha 1 Antibody (SD83-03) (NBP2-67825) is a recombinant monoclonal antibody validated for use in IHC, WB and ICC/IF. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	1291
Gene Symbol	COL6A1
Species	Human, Mouse, Rat
Immunogen	Recombinant protein within Human Collagen VI alpha 1 aa 17-255 / 1,028. (SwissProt: P12109 Human; SwissProt: P12110 Human; SwissProt: P12111 Human; SwissProt: Q02788 Mouse; SwissProt: Q04857 Mouse; Unigene:232118 Rat)


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1:1000-1:2000, Immunohistochemistry, Immunocytochemistry/Immunofluorescence 1:100-1:500, Immunohistochemistry-Paraffin 1:50-1:200

Images


Western Blot: Collagen VI alpha 1 Antibody (SD83-03) [NBP2-67825] - Western blot analysis of Collagen VI alpha 1 on different lysates. Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. The primary antibody (1/500) was used in 5% BSA at room temperature for 2 hours.




Immunocytochemistry/Immunofluorescence: Collagen VI alpha 1 Antibody (SD83-03) [NBP2-67825] - Staining Collagen VI in RH-35 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Immunohistochemistry-Paraffin: Collagen VI alpha 1 Antibody (SD83-03) [NBP2-67825] - Immunofluorescence analysis of paraffin-embedded mouse colon tissue labeling Collagen VI alpha 1 with Rabbit anti-Collagen VI alpha 1 antibody washed with PBS, and then probed with the primary antibody (green) at 1/100 dilution overnight at 4 °C, washed with PBS. Goat Anti-Rabbit IgG H&L (iFluor(TM) 488) was used as the secondary antibody at 1/1,000 dilution. Nuclei were counterstained with DAPI (blue).




Western Blot: Collagen VI alpha 1 Antibody (SD83-03) [NBP2-67825] - Analysis of Collagen VI on different lysates using anti-Collagen VI antibody at 1/1,000 dilution. Positive control: Lane 1: Mouse heart Lane 2: Human heart Lane 3: Human kidney




Immunocytochemistry/Immunofluorescence: Collagen VI alpha 1 Antibody (SD83-03) [NBP2-67825] - Staining Collagen VI in A549 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Immunocytochemistry/Immunofluorescence: Collagen VI alpha 1 Antibody (SD83-03) [NBP2-67825] - Staining Collagen VI in HeLa cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Immunocytochemistry/Immunofluorescence: Collagen VI alpha 1 Antibody (SD83-03) [NBP2-67825] - Staining Collagen VI in HepG2 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Immunohistochemistry-Paraffin: Collagen VI alpha 1 Antibody (SD83-03) [NBP2-67825] - Analysis of paraffin-embedded human liver tissue using anti-Collagen VI antibody. Counter stained with hematoxylin.




Immunohistochemistry-Paraffin: Collagen VI alpha 1 Antibody (SD83-03) [NBP2-67825] - Analysis of paraffin-embedded human lung tissue using anti-Collagen VI antibody. Counter stained with hematoxylin.




Immunohistochemistry-Paraffin: Collagen VI alpha 1 Antibody (SD83-03) [NBP2-67825] - Analysis of paraffin-embedded mouse colon tissue using anti-Collagen VI antibody. Counter stained with hematoxylin.




Immunohistochemistry-Paraffin: Collagen VI alpha 1 Antibody (SD83-03) [NBP2-67825] - Analysis of paraffin-embedded human liver tissue with Rabbit anti-Collagen VI antibody washed with ddH₂O and PBS, and then probed with the primary antibody at 1/400 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Immunohistochemistry-Paraffin: Collagen VI alpha 1 Antibody (SD83-03) [NBP2-67825] - Analysis of paraffin-embedded human colon tissue with Rabbit anti-Collagen VI alpha 1 antibody washed with ddH₂O and PBS, and then probed with the primary antibody at 1/400 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



Immunohistochemistry-Paraffin: Collagen VI alpha 1 Antibody (SD83-03) [NBP2-67825] - Analysis of paraffin-embedded human skin tissue with Rabbit anti-Collagen VI antibody washed with ddH₂O and PBS, and then probed with the primary antibody at 1/400 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



Immunohistochemistry-Paraffin: Collagen VI alpha 1 Antibody (SD83-03) [NBP2-67825] - Immunofluorescence analysis of paraffin-embedded human liver tissue labeling Collagen VI with Rabbit anti-Collagen VI antibody washed with PBS, and then probed with the primary antibody (green) at 1/100 dilution overnight at 4 , washed with PBS. Goat Anti-Rabbit IgG H&L (iFluor(TM) 488) was used as the secondary antibody at 1/1,000 dilution. Nuclei were counterstained with DAPI (blue).





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-67825

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-67825

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

