

Product Datasheet

Cytochrome P450 17A1 Antibody (JB93-32) NBP2-75461

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-75461

Updated 2/24/2026 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-75461



NBP2-75461

Cytochrome P450 17A1 Antibody (JB93-32)


Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	JB93-32
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	TBS (pH7.4), 0.05% BSA, 40% Glycerol
Target Molecular Weight	57 kDa

Product Description	
Description	Novus Biologicals Rabbit Cytochrome P450 17A1 Antibody (JB93-32) (NBP2-75461) is a recombinant monoclonal antibody validated for use in IHC, WB, Flow and ICC/IF. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	1586
Gene Symbol	CYP17A1
Species	Human, Mouse, Rat
Immunogen	Recombinant protein within Human Cytochrome P450 17A1 aa 81-240 / 508. (SwissProt: P05093 Human; SwissProt: P27786 Mouse; SwissProt: P11715 Rat)


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1:500, Flow Cytometry 1:50-1:100, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1:50-1:200, Immunohistochemistry-Paraffin 1:50-1:200

Images


Western Blot: Cytochrome P450 17A1 Antibody (JB93-32) [NBP2-75461]
 - Analysis of Cytochrome P450 17A1 on rat testis tissue lysates with Rabbit anti-Cytochrome P450 17A1 antibody at 1/500 dilution. Lysates/proteins at 20 ug/Lane. Predicted band size: 57 kDa Observed band size: 55 kDa Exposure time: 2 minutes; 10% SDS-PAGE gel. Proteins were transferred to a PVDF membrane and blocked with 5% NFDN/TBST for 1 hour at room temperature. The primary antibody at 1/500 dilution was used in 5% NFDN/TBST at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1:300,000 dilution was used for 1 hour at room temperature.




Immunocytochemistry/Immunofluorescence: Cytochrome P450 17A1 Antibody (JB93-32) [NBP2-75461] - Staining Cytochrome P450 17A1 in SH-SY-5Y cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Immunohistochemistry-Paraffin: Cytochrome P450 17A1 Antibody (JB93-32) [NBP2-75461] - Analysis of paraffin-embedded mouse kidney tissue using anti-Cytochrome P450 17A1 antibody. Counter stained with hematoxylin.




Flow Cytometry: Cytochrome P450 17A1 Antibody (JB93-32) [NBP2-75461] - Analysis of SH-SY-5Y cells with Cytochrome P450 17A1 antibody at 1/100 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti-rabbit IgG was used as the secondary antibody.




Immunocytochemistry/Immunofluorescence: Cytochrome P450 17A1 Antibody (JB93-32) [NBP2-75461] - Staining Cytochrome P450 17A1 in Hela cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Immunocytochemistry/Immunofluorescence: Cytochrome P450 17A1 Antibody (JB93-32) [NBP2-75461] - Staining Cytochrome P450 17A1 in HepG2 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Immunohistochemistry-Paraffin: Cytochrome P450 17A1 Antibody (JB93-32) [NBP2-75461] - Analysis of paraffin-embedded human breast tissue using anti-Cytochrome P450 17A1 antibody. Counter stained with hematoxylin.



Immunohistochemistry-Paraffin: Cytochrome P450 17A1 Antibody (JB93-32) [NBP2-75461] - Analysis of paraffin-embedded human kidney tissue using anti-Cytochrome P450 17A1 antibody. Counter stained with hematoxylin.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-75461

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-75461

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

