

Product Datasheet

TXNIP Antibody (JM60-35)

NBP2-75692

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 4

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-75692

Updated 2/24/2026 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-75692



NBP2-75692

TXNIP Antibody (JM60-35)


Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	JM60-35
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	TBS (pH7.4), 0.05% BSA, 40% Glycerol
Target Molecular Weight	44 kDa

Product Description	
Description	Novus Biologicals Rabbit TXNIP Antibody (JM60-35) (NBP2-75692) is a recombinant monoclonal antibody validated for use in IHC, WB, Flow and ICC/IF. Anti-TXNIP Antibody: Cited in 3 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	10628
Gene Symbol	TXNIP
Species	Human, Mouse, Rat
Immunogen	Synthetic peptide within Human TXNIP aa 50-99 / 391. (SwissProt: Q9H3M7 Human; SwissProt: Q8BG60 Mouse; SwissProt: Q5M7W1 Rat)


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1:500-1:5000, Flow Cytometry 1:50-1:100, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1:50-1:200, Immunohistochemistry-Paraffin 1:50-1:400

Images


Western Blot: TXNIP Antibody (JM60-35) [NBP2-75692] - Analysis of TXNIP on different lysates using anti-TXNIP antibody at 1/1,000 dilution. Positive control: Lane 1: PC-12 Lane 2: Human skeletal muscle Lane 3: NIH-3T3 Lane 4: Hela




Immunocytochemistry/Immunofluorescence: TXNIP Antibody (JM60-35) [NBP2-75692] - Staining TXNIP in NIH-3T3 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Immunohistochemistry-Paraffin: TXNIP Antibody (JM60-35) [NBP2-75692] - Analysis of paraffin-embedded mouse stomach tissue with Rabbit anti-TXNIP antibody washed with ddH₂O and PBS, and then probed with the primary antibody at 1/400 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Flow Cytometry: TXNIP Antibody (JM60-35) [NBP2-75692] - Analysis of HeLa cells with TXNIP antibody at 1/100 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black).




Immunocytochemistry/Immunofluorescence: TXNIP Antibody (JM60-35) [NBP2-75692] - Staining TXNIP in HeLa cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Immunocytochemistry/Immunofluorescence: TXNIP Antibody (JM60-35) [NBP2-75692] - Staining TXNIP in SH-SY5Y cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Immunohistochemistry-Paraffin: TXNIP Antibody (JM60-35) [NBP2-75692] - Analysis of paraffin-embedded human kidney tissue using anti-TXNIP antibody. Counter stained with hematoxylin.




Immunohistochemistry-Paraffin: TXNIP Antibody (JM60-35) [NBP2-75692] - Analysis of paraffin-embedded human stomach cancer tissue using anti-TXNIP antibody. Counter stained with hematoxylin.




Immunohistochemistry-Paraffin: TXNIP Antibody (JM60-35) [NBP2-75692] - Analysis of paraffin-embedded mouse kidney tissue using anti-TXNIP antibody. Counter stained with hematoxylin.




Immunohistochemistry-Paraffin: TXNIP Antibody (JM60-35) [NBP2-75692] - Analysis of paraffin-embedded mouse stomach tissue using anti-TXNIP antibody. Counter stained with hematoxylin.




Immunohistochemistry-Paraffin: TXNIP Antibody (JM60-35) [NBP2-75692] - Analysis of paraffin-embedded human kidney tissue with Rabbit anti-TXNIP antibody washed with ddH₂O and PBS, and then probed with the primary antibody at 1/400 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



Immunohistochemistry-Paraffin: TXNIP Antibody (JM60-35) [NBP2-75692] - Analysis of paraffin-embedded human stomach tissue with Rabbit anti-TXNIP antibody washed with ddH₂O and PBS, and then probed with the primary antibody at 1/400 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



Immunohistochemistry-Paraffin: TXNIP Antibody (JM60-35) [NBP2-75692] - Analysis of paraffin-embedded mouse esophagus tissue with Rabbit anti-TXNIP antibody washed with ddH₂O and PBS, and then probed with the primary antibody at 1/400 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



Publications

Jandova J, Wondrak GT Vemurafenib drives EMT gene expression in BRAFi-resistant BRAFV600E/NRASQ61K melanoma enhancing tumor growth and metastasis in a bioluminescent murine model The Journal of investigative dermatology 2021-10-20 [PMID: 34687745]

Rolver M, Holland L, Prasad N Et al. Chronic Acidosis Rewires Cancer Cell Metabolism Through PPAR alpha Signaling SSRN Electronic Journal 2021-08-18 [PMID: 36533672]

Chu Y, Zhang Y, Gao G, et al. Chorionic Villous Mesenchymal Stem Cells Derived Exosomes Deliver miR-135b-5p to Trophoblasts, Promoting their Proliferation and Invasion by Targeting TXNIP via β -Catenin Pathway Research Square 2020-12-28 (WB)

Jandova J, Wondrak GT Genomic GLO1 deletion modulates TXNIP expression, glucose metabolism, and redox homeostasis while accelerating human A375 malignant melanoma tumor growth Redox biology 2020-12-17 [PMID: 33360689] (IHC-P, WB)





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-75692

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-75692

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

