

Product Datasheet

Ethenoadenosine Antibody (1G4) - Azide and BSA Free NBP2-80727

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-80727

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-80727




NBP2-80727**Ethenoadenosine Antibody (1G4) - Azide and BSA Free**


Product Information	
Unit Size	0.1 ml
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	1G4
Preservative	No Preservative
Isotype	IgG2 Lambda
Purity	Protein G purified
Buffer	PBS
Product Description	
Description	Novus Biologicals Mouse Ethenoadenosine Antibody (1G4) - Azide and BSA Free (NB600-442) is a monoclonal antibody validated for use in WB, ELISA, Flow, ICC/IF and IP. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Species	All Species
Immunogen	Ethenoadenosine
Product Application Details	
Applications	Western Blot, ELISA, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunoprecipitation, CyTOF-ready
Recommended Dilutions	Western Blot 1:800, Flow Cytometry 1:10-1:1000. Use reported in scientific literature (PMID 9774627), ELISA 1:100-1:2000, Immunocytochemistry/Immunofluorescence 1:50-1:100, Immunoprecipitation 1:10-1:500, CyTOF-ready
Application Notes	This antibody is CyTOF ready.

Images


Flow Cytometry: Ethenoadenosine Antibody (1G4) - Azide and BSA Free [NBP2-80727] - Detection of ADP-ribosylated proteins by flow cytometry. C-33A cells (1×10^7 /ml), transiently transfected with a Flag-tagged chicken ART4 containing plasmid, were incubated at 37C in the presence or absence of 200 μ M etheno-NAD⁺ for 30 min. After washing, cells were stained with an etheno-adenosine specific antibody (1G4). Image collected and cropped by CiteAb from the following publication (<https://bmcmolbiol.biomedcentral.com/articles/10.1186/1471-2199-9-86>), licensed under a CC-BY license. Image from the standard format of this antibody.



Flow Cytometry: Ethenoadenosine Antibody (1G4) - Azide and BSA Free [NBP2-80727] - Chicken erythrocytes (1% v/v), isolated by density gradient centrifugation, were transiently transfected with a Flag-tagged chicken ART4 containing plasmid or the empty plasmid (pSecTag B), and incubated at 37C in the presence or absence of 200 μ M etheno-NAD⁺ for 30 min. After washing, cells were stained with an etheno-adenosine specific antibody (1G4) or an anti-Flag antibody and the respective isotype control stained with the 1G4 antibody. Image collected and cropped by CiteAb from the following publication (<https://bmcmolbiol.biomedcentral.com/articles/10.1186/1471-2199-9-86>), licensed under a CC-BY license. Image from the standard format of this antibody.



Flow Cytometry: Ethenoadenosine Antibody (1G4) - Azide and BSA Free [NBP2-80727] - Etheno-ADP-ribosylation of cell-surface proteins on C3H-10T1/2 cells. C3H-10T 1/2 cells at different stages of differentiation (progenitors, MyoD-myoblasts, myotubes) were incubated for 30 minutes in the absence (dashed lines) or presence (solid lines) of 100 μ M etheno-NAD. Cells were then washed, incubated with the primary monoclonal antibody 1G4, stained with a FITC-conjugated goat-anti-mouse secondary antibody and subjected to FACS analysis. Data show one representative experiments out of three. Image collected and cropped by CiteAb from the following publication (<https://bmcmolbiol.biomedcentral.com/articles/10.1186/1471-2199-9-91>), licensed under a CC-BY license. Image from the standard format of this antibody.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-80727

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB7539	Goat anti-Mouse IgG (H+L) Secondary Antibody [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-80727

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

