

Product Datasheet

TEAD1 Antibody - Azide and BSA Free NBP2-95204-0.1ml

Unit Size: 0.1 ml

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-95204

Updated 2/17/2026 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-95204



NBP2-95204-0.1ml

TEAD1 Antibody - Azide and BSA Free


Product Information	
Unit Size	0.1 ml
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Affinity purified
Buffer	PBS (pH 7.3), 50% glycerol
Target Molecular Weight	48 kDa

Product Description	
Description	Novus Biologicals Rabbit TEAD1 Antibody - Azide and BSA Free (NBP2-95204) is a polyclonal antibody validated for use in IHC, WB, ELISA, ICC/IF, IP and ChIP. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	7003
Gene Symbol	TEAD1
Species	Human, Mouse, Rat
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 135-215 of human TEAD1 (NP_068780.2). AIHNKLGLPGIPRPTFFGAPGFWPGMIQTGQPGSSQDVKPFVQQAYPIQPAVT APIPGFEPASAPAPSVPAWQGRSIGTTK


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, ELISA, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunoprecipitation, Chromatin Immunoprecipitation (ChIP)
Recommended Dilutions	Western Blot 1:100 - 1:500, ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements., Immunohistochemistry 1:50 - 1:200, Immunocytochemistry/Immunofluorescence 1:50-1:200, Immunoprecipitation 5µg-4µg antibody for 400µg-600µg extracts of whole cells, Immunohistochemistry-Paraffin 1:50 - 1:200, Chromatin Immunoprecipitation (ChIP) 5µg antibody for 10µg-15µg of Chromatin

Images


Immunocytochemistry/Immunofluorescence: TEF1 Antibody [NBP2-95204] - Analysis of MCF7 cells using TEF1 .



Chromatin Immunoprecipitation: TEF1 Antibody [NBP2-95204] - Analysis of extracts of HeLa cells, using TEF1 rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.



Immunoprecipitation: TEAD1 Antibody - Azide and BSA Free [TEAD1] - Immunoprecipitation of TEAD1 in 600 ug extracts from mouse heart using 3 ug TEAD1 Rabbit pAb . Western blot analysis was performed using TEAD1 Rabbit pAb at 1:200 dilution.



Western Blot: TEAD1 Antibody - Azide and BSA Free [TEAD1] - Western blot analysis of lysates from Rat lung using TEAD1 Rabbit pAb at 1:2000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1:10000 dilution.

Lysates/proteins: 25 ug per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit .


Exposure time:90s.



Immunohistochemistry: TEAD1 Antibody - Azide and BSA Free [TEAD1]
 - Immunohistochemistry analysis of paraffin-embedded Human liver using TEAD1 Rabbit pAb at dilution of 1:50 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Immunohistochemistry: TEAD1 Antibody - Azide and BSA Free [TEAD1]
 - Immunohistochemistry analysis of paraffin-embedded Human lung using TEAD1 Rabbit pAb at dilution of 1:50 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Western Blot: TEAD1 Antibody - Azide and BSA Free [TEAD1] - Western blot analysis of various lysates using TEAD1 Rabbit pAb at 1:700 dilution incubated overnight at 4C.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1:10000 dilution.

Lysates/proteins: 25 ug per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit .

Exposure time: 28s.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-95204-0.1ml

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-95204

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

