

Product Datasheet

IFI16 Antibody - BSA Free NBP3-18336-0.1mg

Unit Size: 0.1 mg

Store at 4C.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP3-18336

Updated 10/13/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP3-18336



NBP3-18336-0.1mg

IFI16 Antibody - BSA Free


Product Information	
Unit Size	0.1 mg
Concentration	1 mg/ml
Storage	Store at 4C.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Peptide affinity purified
Buffer	PBS

Product Description	
Description	Novus Biologicals Rabbit IFI16 Antibody - BSA Free (NBP3-18336) is a polyclonal antibody validated for use in IHC, WB, ELISA and ICC/IF. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	3428
Gene Symbol	IFI16
Species	Human, Mouse, Rat
Immunogen	Antibody was raised against a peptide corresponding to 16 amino acids near the carboxyl terminus of human IFI16. The immunogen is located within the last 50 amino acids of IFI16.


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, ELISA, Immunocytochemistry/Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 2 ug/mL, ELISA, Immunohistochemistry 1 ug/ml, Immunocytochemistry/ Immunofluorescence 20 ug/mL, Immunohistochemistry-Paraffin

Images


Western Blot: IFI16 Antibody [NBP3-18336] - Western Blot Validation in Rat Testis. Loading: 15 ug of lysate per lane. Antibodies: IFI16 NBP3-18336, 2 ug/mL, 1h incubation at RT in 5% NFDm/TBST. Secondary: Goat anti-rabbit IgG HRP conjugate at 1:10,000 dilution.




Western Blot: IFI16 Antibody [NBP3-18336] - WB Validation in Human and Rat Cell Lines. Loading: 15 ug of lysate per lane. Antibodies: IFI16 NBP3-18336, 2 ug/mL, 1 h incubation at RT in 5% NFDN/TBST. Secondary: Goat Anti-Rabbit IgG HRP conjugate at 1:10,000 dilution.




Western Blot: IFI16 Antibody [NBP3-18336] - Western Blot Validation in Mouse Tissues. Loading: 15 ug of lysates per lane. Antibodies: IFI16 NBP3-18336, 2 ug/mL, 1h incubation at RT in 5% NFDN/TBST. Secondary: Goat anti-rabbit IgG HRP conjugate at 1:10,000 dilution.




Immunohistochemistry-Paraffin: IFI16 Antibody [NBP3-18336] - Immunohistochemical analysis of paraffin-embedded human testis tissue using anti-IFI16 antibody (NBP3-18336) at 1 ug/mL. Tissue was fixed with formaldehyde and blocked with 10% serum for 1 h at RT; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody overnight at 4C. A goat anti-rabbit IgG H&L (HRP) at 1/250 was used as secondary. Counter stained with Hematoxylin.




Immunohistochemistry-Paraffin: IFI16 Antibody [NBP3-18336] - Immunohistochemical analysis of paraffin-embedded mouse liver tissue using anti-IFI16 antibody (NBP3-18336) at 1 ug/mL. Tissue was fixed with formaldehyde and blocked with 10% serum for 1 h at RT; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody overnight at 4C. A goat anti-rabbit IgG H&L (HRP) at 1/250 was used as secondary. Counter stained with Hematoxylin.



Immunocytochemistry/ Immunofluorescence: IFI16 Antibody - BSA Free [NBP3-18336] - Validation of IFI16 in Human Molt4 Cells. Immunofluorescent analysis of 4% paraformaldehyde-fixed Molt4 cells labeling IFI16 with at 20 ug/mL, followed by goat anti-rabbit IgG secondary antibody at 1/500 dilution (green) and DAPI staining (blue).



Immunohistochemistry: IFI16 Antibody - BSA Free [NBP3-18336] - Validation of IFI16 in Rat Testis. Immunohistochemical analysis of paraffin-embedded rat testis tissue using anti-IFI16 antibody at 1 ug/mL. Tissue was fixed with formaldehyde and blocked with 10% serum for 1 h at RT; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody overnight at 4C. A goat anti-rabbit IgG H&L (HRP) at 1/250 was used as secondary. Counter stained with Hematoxylin.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP3-18336-0.1mg

NBP3-18336PEP	IF116 Antibody Blocking Peptide
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-18336

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

