

Product Datasheet

CD19 Antibody (PSH03-94)

NBP3-32130

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP3-32130

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP3-32130



NBP3-32130

CD19 Antibody (PSH03-94)

Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	PSH03-94
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	PBS (pH7.4), 0.1% BSA and 40% Glycerol
Target Molecular Weight	60 kDa

Product Description	
Description	Novus Biologicals Rabbit CD19 Antibody (PSH03-94) (NBP3-32130) is a recombinant monoclonal antibody validated for use in IHC, WB and IP. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	930
Gene Symbol	CD19
Species	Mouse
Immunogen	Recombinant protein within mouse CD19 protein. (Uniprot: P25918)


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Immunoprecipitation
Recommended Dilutions	Western Blot 1:1000, Immunohistochemistry, Immunoprecipitation 1-2ug/sample, Immunohistochemistry-Paraffin 1:500

Images

Immunoprecipitation: CD19 Antibody (PSH03-94) [NBP3-32130] - CD19 was immunoprecipitated in 0.2mg Mouse spleen tissue lysate with NBP3-32130 at 2 ug/25 ul agarose. Western blot was performed from the immunoprecipitate using NBP3-32130 at 1/1,000 dilution. Anti-Rabbit IgG for IP Nano-secondary antibody at 1/5,000 dilution was used for 1 hour at room temperature.

Lane 1: Mouse spleen tissue lysate (input)
 Lane 2: NBP3-32130 IP in Mouse spleen tissue lysate
 Lane 3: Rabbit IgG instead of NBP3-32130 in Mouse spleen tissue lysate

Blocking/Dilution buffer: 5% NFDM/TBST
 Exposure time: 17 seconds



Western Blot: CD19 Antibody (PSH03-94) [NBP3-32130] - Western blot analysis of CD19 on mouse spleen tissue lysates with Rabbit anti-CD19 antibody (NBP3-32130) at 1/1,000 dilution.

Lysates/proteins at 20 ug/Lane.


Predicted band size: 60 kDa

Observed band size: 60-120 kDa

Exposure time: 30 seconds;


4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (NBP3-32130) at 1/1,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1/50,000 dilution was used for 1 hour at room temperature.




Immunohistochemistry: CD19 Antibody (PSH03-94) [NBP3-32130] - Immunofluorescence analysis of paraffin-embedded mouse spleen tissue labeling CD19 with Rabbit anti-CD19 antibody (NBP3-32130) at 1/200 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 10% negative goat serum for 1 hour at room temperature, washed with PBS, and then probed with the primary antibody (NBP3-32130, green) at 1/200 dilution overnight at 4 °C, washed with PBS. Goat Anti-Rabbit IgG H&L (iFluor™ 488) was used as the secondary antibody at 1/1,000 dilution. Nuclei were counterstained with DAPI (blue).



Immunohistochemistry: CD19 Antibody (PSH03-94) [NBP3-32130] - Immunohistochemical analysis of paraffin-embedded mouse lymph node tissue with Rabbit anti-CD19 antibody (NBP3-32130) at 1/500 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (NBP3-32130) at 1/500 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP3-32130

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-32130

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

