

Product Datasheet

NDUFV2 Antibody (PSH02-86) NBP3-32640

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP3-32640

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP3-32640



NBP3-32640

NDUFV2 Antibody (PSH02-86)

Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	PSH02-86
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	PBS (pH7.4), 0.1% BSA and 40% Glycerol
Target Molecular Weight	27 kDa
Product Description	
Description	Novus Biologicals Rabbit NDUFV2 Antibody (PSH02-86) (NBP3-32640) is a recombinant monoclonal antibody validated for use in IHC, WB and ICC/IF. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	4729
Gene Symbol	NDUFV2
Species	Human, Mouse, Rat
Immunogen	Recombinant protein within human NDUFV2 aa 1-249 / 249. (Uniprot: P19404)
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1:2000, Immunohistochemistry, Immunocytochemistry/Immunofluorescence 1:2000, Immunohistochemistry-Paraffin 1:2000-1:5000


Images

Immunohistochemistry: NDUFV2 Antibody (PSH02-86) [NBP3-32640] - Immunohistochemical analysis of paraffin-embedded human heart tissue with Rabbit anti-NDUFV2 antibody (NBP3-32640) at 1/2,000 dilution. ✖

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody at 1/2,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Immunocytochemistry/ Immunofluorescence: NDUFV2 Antibody (PSH02-86) [NBP3-32640] - Immunocytochemistry analysis of RAW264.7 cells labeling NDUFV2 with Rabbit anti-NDUFV2 antibody (NBP3-32640) at 1/2,000 dilution.

Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-NDUFV2 antibody at 1/2,000 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.



Western Blot: NDUFV2 Antibody (PSH02-86) [NBP3-32640] - Western blot analysis of NDUFV2 on different lysates with Rabbit anti-NDUFV2 antibody (NBP3-32640) at 1/2,000 dilution.


Lane 1: A549 cell lysate (20 µg/Lane)
 Lane 2: HeLa cell lysate (20 µg/Lane)
 Lane 3: Jurkat cell lysate (20 µg/Lane)
 Lane 4: Ramos cell lysate (20 µg/Lane)
 Lane 5: Raji cell lysate (20 µg/Lane)
 Lane 6: K-562 cell lysate (20 µg/Lane)
 Lane 7: A431 cell lysate (20 µg/Lane)
 Lane 8: NIH/3T3 cell lysate (20 µg/Lane)
 Lane 9: RAW264.7 cell lysate (20 µg/Lane)
 Lane 10: PC-12 cell lysate (20 µg/Lane)
 Lane 11: Mouse heart tissue lysate (40 µg/Lane)
 Lane 12: Rat heart tissue lysate (40 µg/Lane)
 Lane 13: Human kidney tissue lysate (40 µg/Lane)
 Lane 14: Mouse kidney tissue lysate (40 µg/Lane)

Predicted band size: 27 kDa
 Observed band size: 24 kDa

Exposure time: 1 minute 59 seconds;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDN/TBST for 1 hour at room temperature. The primary antibody at 1/2,000 dilution was used in 5% NFDN/TBST at 4 °C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1/50,000 dilution was used for 1 hour at room temperature.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA

Phone: 303.730.1950

Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada

Phone: 905.827.6400

Toll Free: 855.668.8722

Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15

Fax: (44) (0) 1235 533420

info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-32640

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-32640

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

