

Product Datasheet

Rad17 [p Ser656] Antibody (JE63-30) NBP3-32792

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP3-32792

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP3-32792



NBP3-32792

Rad17 [p Ser656] Antibody (JE63-30)

Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	JE63-30
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	1*TBS (pH7.4), 0.05% BSA and 40% Glycerol
Target Molecular Weight	77 kDa

Product Description	
Description	Novus Biologicals Rabbit Rad17 [p Ser656] Antibody (JE63-30) (NBP3-32792) is a recombinant monoclonal antibody validated for use in WB, Flow and ICC/IF. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	5884
Gene Symbol	RAD17
Species	Human, Mouse
Immunogen	Synthetic peptide corresponding to residues surrounding Ser656 of human Rad17 protein. (Uniprot: O75943)

Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence
Recommended Dilutions	Western Blot 1:1000, Flow Cytometry 1:1000, Immunocytochemistry/ Immunofluorescence 1:500



Images

Western Blot: Rad17 [p Ser656] Antibody (JE63-30) [NBP3-32792] - Western blot analysis of Rad17 on different lysates with Rabbit anti-Rad17 antibody (NBP3-32792) at 1/1,000 dilution.

Lane 1: HeLa cell lysate

Lane 2: HeLa treated with 100ng/mL Calyculin A for 30 minutes cell lysate

Lane 3: HeLa treated with 100ng/mL Calyculin A for 30 minutes cell lysate, then the membrane treated with pp for 1 hour

Lane 4: NIH/3T3 cell lysate

Lane 5: NIH/3T3 treated with 50ng/mL Calyculin A for 45 minutes cell lysate

Lane 6: NIH/3T3 treated with 50ng/mL Calyculin A for 45 minutes cell lysate, then the membrane treated with pp for 1 hour

Lysates/proteins at 20 ug/Lane.


Predicted band size: 77 kDa

Observed band size: 77 kDa

Exposure time: 43 seconds; ECL;


4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (NBP3-32792) at 1/1,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1/50,000 dilution was used for 1 hour at room temperature.




Immunocytochemistry/ Immunofluorescence: Rad17 [p Ser656] Antibody (JE63-30) [NBP3-32792] - Immunocytochemistry analysis of HeLa cells treated with or without 100ng/mL Calyculin A for 30 minutes labeling Rad17 with Rabbit anti-Rad17 antibody (NBP3-32792) at 1/500 dilution.

Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-Rad17 antibody (NBP3-32792) at 1/500 dilution in 1% BSA in PBST overnight at 4°C. Goat Anti-Rabbit IgG H&L (iFluor™ 488) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.



Flow Cytometry: Rad17 [p Ser656] Antibody (JE63-30) [NBP3-32792] - Flow cytometric analysis of HeLa cells treated with or without 100ng/mL Calyculin A for 30 minutes labeling Rad17.

Cells were fixed and permeabilized. Then stained with the primary antibody (NBP3-32792, 1 μ g/mL) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4°C for an hour, the cells were stained with a iFluor™ 488 conjugate-Goat anti-Rabbit IgG Secondary antibody at 1/1,000 dilution for 30 minutes at +4°C. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA

Phone: 303.730.1950

Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada

Phone: 905.827.6400

Toll Free: 855.668.8722

Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15

Fax: (44) (0) 1235 533420

info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-32792

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-32792

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

