

Product Datasheet

PPP2R5D Antibody (PSH01-21) NBP3-32846

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP3-32846

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP3-32846



NBP3-32846

PPP2R5D Antibody (PSH01-21)

Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	PSH01-21
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	PBS (pH7.4), 0.1% BSA and 40% Glycerol
Target Molecular Weight	70 kDa

Product Description	
Description	Novus Biologicals Rabbit PPP2R5D Antibody (PSH01-21) (NBP3-32846) is a recombinant monoclonal antibody validated for use in WB, Flow and ICC/IF. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	5528
Gene Symbol	PPP2R5D
Species	Human
Immunogen	Recombinant protein within human PPP2R5D aa 107-602 / 602. (Uniprot: Q14738)

Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence
Recommended Dilutions	Western Blot 1:1000, Flow Cytometry 1:500-1:1000, Immunocytochemistry/ Immunofluorescence 1:100



Images

Western Blot: PPP2R5D Antibody (PSH01-21) [NBP3-32846] - Western blot analysis of PPP2R5D on different lysates with Rabbit anti-PPP2R5D antibody (NBP3-32846) at 1/1,000 dilution.

Lane 1: A431 cell lysate (20 ug/Lane)
 Lane 2: HeLa cell lysate (20 ug/Lane)
 Lane 3: HepG2 cell lysate (20 ug/Lane)
 Lane 4: MCF7 cell lysate (20 ug/Lane)
 Lane 5: 293T cell lysate (20 ug/Lane)
 Lane 6: Jurkat cell lysate (20 ug/Lane)
 Lane 7: Human brain tissue lysate (40 ug/Lane)

Predicted band size: 70 kDa
 Observed band size: 56~70 kDa

Exposure time: 2 minutes 10 seconds;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDN/TBST for 1 hour at room temperature. The primary antibody (NBP3-32846) at 1/1,000 dilution was used in 5% NFDN/TBST at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1:100,000 dilution was used for 1 hour at room temperature.




Immunocytochemistry/ Immunofluorescence: PPP2R5D Antibody (PSH01-21) [NBP3-32846] - Immunocytochemistry analysis of A431 cells labeling PPP2R5D with Rabbit anti-PPP2R5D antibody (NBP3-32846) at 1/100 dilution.

Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-PPP2R5D antibody (NBP3-32846) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (red) was stained at 1/100 dilution overnight at +4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 594) was used as the secondary antibody at 1/1,000 dilution.

Flow Cytometry: PPP2R5D Antibody (PSH01-21) [NBP3-32846] - Flow cytometric analysis of A431 cells labeling PPP2R5D.

Cells were fixed and permeabilized. Then stained with the primary antibody (NBP3-32846, 1ug/ml) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4 °C for an hour, the cells were stained with a iFluor™ 488 conjugate-Goat anti-Rabbit IgG Secondary antibody at 1/1,000 dilution for 30 minutes at +4 °C. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP3-32846

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-32846

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

