

# Product Datasheet

## FPRL1/FPR2 Antibody - BSA Free NLS1878

Unit Size: 0.05 ml

Store at -20 degrees C. Avoid freeze/thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

**Reviews: 4 Publications: 42**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NLS1878](http://www.novusbio.com/NLS1878)

Updated 4/21/2026 v.20.1

**Earn rewards for product  
reviews and publications.**

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NLS1878](http://www.novusbio.com/reviews/destination/NLS1878)



**NLS1878**

FPRL1/FPR2 Antibody - BSA Free

Product Information	
Unit Size	0.05 ml
Concentration	1.1 mg/ml
Storage	Store at -20 degrees C. Avoid freeze/thaw cycles.
Clonality	Polyclonal
Preservative	0.01% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS, 30% Glycerol
Target Molecular Weight	38 kDa


Product Description	
Host	Rabbit
Gene ID	2358
Gene Symbol	FPR2
Species	Human, Mouse, Rat, Bacteria
Reactivity Notes	Rat reactivity reported in scientific literature (PMID: 24086560). Bacteria reactivity reported in scientific literature (PMID: 31234710).
Immunogen	This FPRL1/FPR2 antibody is made to a synthetic peptide from the human FPRL1 protein (between residues 300-350) [UniProt P25090]

Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Electron Microscopy, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen
Recommended Dilutions	Western Blot 1:1000, Flow Cytometry, Immunohistochemistry 1:300, Immunocytochemistry/ Immunofluorescence 1:20-1:75, Immunohistochemistry-Paraffin 1:300, Immunohistochemistry-Frozen reported in scientific literature (PMID 24086560), Electron Microscopy
Application Notes	In Western Blot, a band is seen ~38 kDa representing FPRL1. In ICC/IF, plasma membrane staining was observed in Raw264.7 cells. In IHC-P, staining was also observed in the plasma membrane of human kidney cancer tissue. Prior to immunostaining paraffin tissues, antigen retrieval with sodium citrate buffer (pH 6.0) is recommended. Customers have reported success in IF on FFPE mouse kidney tissue, following microwave antigen retrieval.




## Images


**Western Blot: FPRL1/FPR2 Antibody - BSA Free [NLS1878]** - Immunoblot of FPR2 protein in various parts of the rat brain including olfactory bulb (OB), prefrontal cortex (PFC), primary somatosensory cortex (CTX1), parietal association cortex and secondary auditory cortex (CTX2), striatum (STR), thalamus and hypothalamus (THA), hippocampus (HPC), cerebellum (CB), brainstem (BS), cervical spinal cord [SC(C)], thoracic spinal cord [SC(T)], and lumbar spinal cord [SC(L)]. Blots incubated with antigen-absorbed antibody i. e. peptide competition, show reduced band intensities. Image collected and cropped by Citeab from the following publication (Localisation of Formyl-Peptide Receptor 2 in the Rat Central Nervous System and Its Role in Axonal and Dendritic Outgrowth. *Neurochem Res* (2018)) licensed under a CC-BY license.




**Immunocytochemistry/Immunofluorescence: FPRL1/FPR2 Antibody - BSA Free [NLS1878]** - Mouse J774A.1, mouse reticulum cell sarcoma macrophage cell line. Left panel is + Anti-Rabbit FITC; middle panel indicates the cell nuclei stained with Hoechst; the right panel is a merged image. ICC/IF image submitted by a verified customer review.




**Immunohistochemistry-Paraffin: FPRL1/FPR2 Antibody - BSA Free [NLS1878]** - SPM receptors differentially expressed on endothelium and infiltrating leukocytes AE" ALX/FPR2 and ChemR23. Acute inflammation triggered by ventral aspect of forearm of healthy volunteers by intradermal injection of 1.5 A-107 UV-killed E.coli (UVkEc) suspended in 100  $\mu$ l of saline. 4hrs after injection a 3-mm skin punch biopsy taken from inflamed site under local anesthesia. Naive skin treated as baseline. IHC-P on skin sections for receptor identification. Low mag(A-5) and high-mag (A-40) images at baseline and the 4 hr time point shown for ALX/FPR2. Red arrows highlight endothelium. Black arrow highlights infiltrating leukocytes. Image collected and cropped by Citeab from the (Pro-resolving mediators promote resolution in human skin model of UV-killed Escherichia coli-driven acute inflammation *JCI Insight* (2018)) licensed under, CC-BY license.



**Flow Cytometry: FPRL1/FPR2 Antibody - BSA Free [NLS1878]** - Flow Cytometry: [Alexa Fluor® 700] [NLS1878AF700] - Mouse splenocytes. Cells were pre-gated with live/dead and FSC-A/W to exclude dead cells and cell doublets. Image from verified customer review. Image using the Alexa Fluor 700 form of this antibody.




Western Blot: FPRL1/FPR2 Antibody - BSA Free [NLS1878] - Murine neutrophils were extracted from the bone marrow of C57BL/6 mice and exposed to normal glucose (90 mg/dl) or high glucose (300 mg/dl) and the expression of FPR2 was assessed by western blotting after 1, 2, or 3 hr exposure to glucose. Image collected and cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/35112667/>) licensed under a CC-BY license.




Western Blot: FPRL1/FPR2 Antibody - BSA Free [NLS1878] - Analysis in HL-60 cell lysate.




Immunocytochemistry/Immunofluorescence: FPRL1/FPR2 Antibody - BSA Free [NLS1878] - Antibody was tested in Raw264.7 cells with DyLight 488 (green). Nuclei were counterstained with DAPI (blue).




Immunohistochemistry: FPRL1/FPR2 Antibody - BSA Free [NLS1878] - Analysis in human kidney cancer using DAB with hematoxylin counterstain.




**Electron Microscopy: FPRL1/FPR2 Antibody - BSA Free [NLS1878] -**  
 Electron micrographs of FPR2 immunostained sections from the prefrontal cortex. Immunostaining is mostly present in axon pre-terminals (PrT) that did not form synapses with postsynaptic structures. Occasional labelled dendrites (DE) are also found, that formed asymmetrical synapses (S) with unlabelled axon terminals (AT). Scale: a, b, c=50A nm, e=100A nm. Image collected and cropped by Citeab from the following publication (Localisation of Formyl-Peptide Receptor 2 in the Rat Central Nervous System and Its Role in Axonal and Dendritic Outgrowth. *Neurochem Res* (2018)) licensed under a CC-BY license.




**Immunohistochemistry: FPRL1/FPR2 Antibody - BSA Free [NLS1878] -**  
 Electron micrographs of FPR2 immunostained sections from the prefrontal cortex. a, b Immunostaining is mostly present in axon pre-terminals (PrT) that did not form synapses with postsynaptic structures. c Occasional axon terminals (AT) are observed to form asymmetrical, putatively glutamatergic synapses (S) with unlabelled dendrites (DE). d Occasional labelled dendrites (DE) are also found, that formed asymmetrical synapses (S) with unlabelled axon terminals (AT). Scale: a, b, c = 50 nm, e = 100 nm Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/29948727>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




**Immunohistochemistry: FPRL1/FPR2 Antibody - BSA Free [NLS1878] -**  
 Electron micrographs of FPR2 immunostained sections from the prefrontal cortex. a, b Immunostaining is mostly present in axon pre-terminals (PrT) that did not form synapses with postsynaptic structures. c Occasional axon terminals (AT) are observed to form asymmetrical, putatively glutamatergic synapses (S) with unlabelled dendrites (DE). d Occasional labelled dendrites (DE) are also found, that formed asymmetrical synapses (S) with unlabelled axon terminals (AT). Scale: a, b, c = 50 nm, e = 100 nm Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/29948727>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




**Immunohistochemistry: FPRL1/FPR2 Antibody - BSA Free [NLS1878] -**  
 Electron micrographs of FPR2 immunostained sections from the prefrontal cortex. a, b Immunostaining is mostly present in axon pre-terminals (PrT) that did not form synapses with postsynaptic structures. c Occasional axon terminals (AT) are observed to form asymmetrical, putatively glutamatergic synapses (S) with unlabelled dendrites (DE). d Occasional labelled dendrites (DE) are also found, that formed asymmetrical synapses (S) with unlabelled axon terminals (AT). Scale: a, b, c = 50 nm, e = 100 nm Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/29948727>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




**Immunohistochemistry: FPRL1/FPR2 Antibody - BSA Free [NLS1878] -** Electron micrographs of FPR2 immunostained sections from the prefrontal cortex. a, b Immunostaining is mostly present in axon pre-terminals (PrT) that did not form synapses with postsynaptic structures. c Occasional axon terminals (AT) are observed to form asymmetrical, putatively glutamatergic synapses (S) with unlabelled dendrites (DE). d Occasional labelled dendrites (DE) are also found, that formed asymmetrical synapses (S) with unlabelled axon terminals (AT). Scale: a, b, c = 50 nm, e = 100 nm Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/29948727>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



**Immunocytochemistry/ Immunofluorescence: FPRL1/FPR2 Antibody - BSA Free [NLS1878] -** Lipoxin A4, along with its precursors & ligand, were altered by hAECs. (A): Expression of the lipoxin A4 precursor genes ALOX $\zeta$ 5,  $\zeta$ 12, &  $\zeta$ 15 was increased at days 5 & 7 compared with saline controls (ALOX $\zeta$ 5:  $0.393 \pm 0.114$  vs.  $3.777 \pm 1.98$ ; ALOX $\zeta$ 12:  $0.484 \pm 0.248$  vs.  $1.520 \pm 0.203$ ; ALOX $\zeta$ 15:  $0.020 \pm 0.007$  vs.  $1.998 \pm 0.983$ ).  $\square$ ,  $p < .05$ . (B): Lipoxin A4 protein levels were elevated in lung lysates at day 7 in animals treated with hAECs compared with controls ( $0.103 \pm 0.021$  ng/ml vs.  $0.249 \pm 0.072$  ng/ml, respectively).  $\square$ ,  $p < .05$ . (C): At day 7, bleomycin challenge resulted in positively stained F4/80/FPR2 cells, which was elevated in mice treated with hAECs compared with control mice ( $5.720\% \pm 0.587\%$  vs.  $8.795\% \pm 0.687\%$ , respectively).  $\square$ ,  $p < .05$ . (D): Representative images of F4/80 $\square$  & FPR2 $\square$ positive $\square$ stained lung sections from hAEC $\square$ treated animals. Magnification:  $\times 200$ . Scale bar =  $100 \mu\text{m}$ . Abbreviations: DAPI, 4',6-diamidino-2-phenylindole; FPR2, N-formyl peptide receptor 2; hAEC, human amnion epithelial cell. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/28371562>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



**Western Blot: FPRL1/FPR2 Antibody - BSA Free [NLS1878] -** Higher expression of Fpr2 in the livers of female mice is related with hepatocyte protection. a qRT-PCR analysis for Fpr2 expression in primary hepatocytes (pHEPs) from WT (male  $n = 2$ , female  $n = 2$ ) & KO mice (male  $n = 2$ , female  $n = 2$ ). Total eight mice were employed in each hepatocyte isolation, & the experiments were replicated at least three times & the mean  $\pm$  S.E.M. results are graphed ( $*p < 0.05$ ,  $**p < 0.005$  vs WT male-pHEPs). b Western blot analysis & c double immunofluorescent images of Fpr2 (red) with albumin (green) in these cells. Gapdh was used as internal control. DAPI (blue) was used as nuclear counterstaining. Data shown represent one of three experiments with similar results (Scale bar,  $20 \mu\text{m}$ ). d qRT-PCR analysis for Fpr2 & glucose-6-phosphatase (G6pc) in, e cell viability of, & f western blot analysis of cleaved Caspase-3 & pro Caspase-3 in WT & KO female mice-isolated pHEPs treated with vehicle (Veh) or  $250 \mu\text{M}$  of palmitate (PA). The data shown represent one of three experiments with similar results. The mean  $\pm$  S.E.M. results obtained from three repetitive experiments are graphed ( $*p < 0.05$ ,  $**p < 0.005$  vs WT-Veh). Gray circles represent individual data points. See Supplementary Data for statistical details. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/35102146>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



## Publications

Uno H, Itokazu T, Yamashita T Formyl peptide receptor 2 antagonist WRW4 ameliorates diabetes-induced cognitive decline in mice. *Neuroscience Research* 2025-07-05 [PMID: 40618851]

Zhang Q, Zha Y, Wang X et al. FPR2 Agonism Attenuates Restenosis by Mitigating Neointimal Hyperplasia via ELOVL6 *The FASEB Journal* 2025-09-06 [PMID: 40913417]

Li K, Yi Y, Ling R et al. PCIF1 drives oesophageal squamous cell carcinoma progression via m6Am-mediated suppression of MTF2 translation. *Clinical and translational medicine* 2025-03-29 [PMID: 40156159]

Lin T, Cui Z, Shen Y et al. Elevated LL-37/FPR2 Axis and its Regulatory Role for Gingival Fibroblasts in Periodontitis. *International dental journal* 2025-10-09 [PMID: 41072077]

YE Kim, SY Ahn, DK Sung, YS Chang, WS Park Mesenchymal Stem Cells and Formyl Peptide Receptor 2 Activity in Hyperoxia-Induced Lung Injury in Newborn Mice *International Journal of Molecular Sciences*, 2022-09-13;23(18):. 2022-09-13 [PMID: 36142517] (Western Blot, Human, Mouse)

R Roy, J Zayas, SK Singh, K Delgado, SJ Wood, MF Mohamed, DM Frausto, YA Albalawi, TP Price, R Estupinian, EF Giurini, TM Kuzel, A Zloza, J Reiser, SH Shafikhani Overriding impaired FPR chemotaxis signaling in diabetic neutrophil stimulates infection control in murine diabetic wound *Elife*, 2022-02-03;11(0):. 2022-02-03 [PMID: 35112667] (Western Blot, Human, Mouse)

C Lee, J Kim, J Han, D Oh, M Kim, H Jeong, TJ Kim, SW Kim, JN Kim, YS Seo, A Suzuki, JH Kim, Y Jung Formyl peptide receptor 2 determines sex-specific differences in the progression of nonalcoholic fatty liver disease and steatohepatitis *Nature Communications*, 2022-01-31;13(1):578. 2022-01-31 [PMID: 35102146] (Western Blot, Human, Mouse)

Xie T, Cai J, Yao Y et al. LXA4 protects against blue-light induced retinal degeneration in human A2E-laden RPE cells and Balb-c mice *Annals of Translational Medicine* 2021-08-01 [PMID: 34532386] (Western Blot, Human, Mouse)

Zamani A, EmamiAref P, Kubí?ková L, Haanová K et Al. Paclitaxel triggers molecular and cellular changes in the choroid plexus *Front Pain Res (Lausanne)* 2024-12-10 [PMID: 39654799]

Yang SC, Wang YH, Ho CM et al. Targeting formyl peptide receptor 1 by anteiso-C13-surfactin for neutrophil-dominant acute respiratory distress syndrome *British journal of pharmacology* 2023-03-16 [PMID: 36929355]

Zhu M, Wang Y, Zhu L et al. Crosstalk between RPE cells and choroidal endothelial cells via the ANXA1/FPR2/SHP2/NLRP3 inflammasome/pyroptosis axis promotes choroidal neovascularization *Inflammation* 2021-10-01 [PMID: 34595678]

Federica Frigerio, Giulia Pasqualini, Ilaria Craparotta, Sergio Marchini, Erwin A van Vliet, Patrick Foerch, Catherine Vandenplas, Karin Leclercq, Eleonora Aronica, Luca Porcu, Kimberly Pistorius, Romain A Colas, Trond V Hansen, Mauro Perretti, Rafal M Kaminski, Jesmond Dalli, Annamaria Vezzani n-3 Docosapentaenoic acid-derived protectin D1 promotes resolution of neuroinflammation and arrests epileptogenesis *Brain* 2018-11-01 [PMID: 30307467]

More publications at <http://www.novusbio.com/NLS1878>



## Procedures

### Immunohistochemistry-Paraffin protocol for FPRL1 Antibody (NLS1878)

#### Immunohistochemistry-Paraffin Embedded Sections

##### Antigen Unmasking:

Bring slides to a boil in 10 mM sodium citrate buffer (pH 6.0) then maintain at a sub-boiling temperature for 10 minutes. Cool slides on bench-top for 30 minutes.

##### Staining:

1. Wash sections in deionized water three times for 5 minutes each.
2. Wash sections in wash buffer for 5 minutes.
3. Block each section with 100-400 ul blocking solution for 1 hour at room temperature.
4. Remove blocking solution and add 100-400 ul diluted primary antibody. Incubate overnight at 4 C.
5. Remove antibody solution and wash sections in wash buffer three times for 5 minutes each.
6. Add 100-400 ul biotinylated diluted secondary antibody. Incubate 30 minutes at room temperature.
7. Remove secondary antibody solution and wash sections three times with wash buffer for 5 minutes each.
8. Add 100-400 ul Streptavidin-HRP reagent to each section and incubate for 30 minutes at room temperature.
9. Wash sections three times in wash buffer for 5 minutes each.
10. Add 100-400 ul DAB substrate to each section and monitor staining closely.
11. As soon as the sections develop, immerse slides in deionized water.
12. Counterstain sections in hematoxylin.
13. Wash sections in deionized water two times for 5 minutes each.
14. Dehydrate sections.
15. Mount coverslips.

\*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.

### Western Blot protocol for FPRL1 Antibody (NLS1878)

#### Western Blot Protocol

1. Perform SDS-PAGE on samples to be analyzed, loading 40 ug of total protein per lane.
2. Transfer proteins to membrane according to the instructions provided by the manufacturer of the membrane and transfer apparatus.
3. Stain according to standard Ponceau S procedure (or similar product) to assess transfer success, and mark molecular weight standards where appropriate.
4. Rinse the blot.
5. Block the membrane using standard blocking buffer for at least 1 hour.
6. Wash the membrane in wash buffer three times for 10 minutes each.
7. Dilute primary antibody in blocking buffer and incubate 1 hour at room temperature.
8. Wash the membrane in wash buffer three times for 10 minutes each.
9. Apply the diluted HRP conjugated secondary antibody in blocking buffer (as per manufacturers instructions) and incubate 1 hour at room temperature.
10. Wash the blot in wash buffer three times for 10 minutes each (this step can be repeated as required to reduce background).
11. Apply the detection reagent of choice in accordance with the manufacturers instructions.

Note: Tween-20 can be added to the blocking or antibody dilution buffer at a final concentration of 0.05-0.2%.

\*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.

**Immunocytochemistry/Immunofluorescence Protocol for FPRL1 Antibody (NLS1878)**

## Immunocytochemistry Protocol

Culture cells to appropriate density in 35 mm culture dishes or 6-well plates.

1. Remove culture medium and wash the cells briefly in PBS. Add 10% formalin to the dish and fix at room temperature for 10 minutes.
2. Remove the formalin and wash the cells in PBS.
3. Permeabilize the cells with 0.1% Triton X100 or other suitable detergent for 10 min.
4. Remove the permeabilization buffer and wash three times for 10 minutes each in PBS. Be sure to not let the specimen dry out.
5. To block nonspecific antibody binding, incubate in 10% normal goat serum from 1 hour to overnight at room temperature.
6. Add primary antibody at appropriate dilution and incubate overnight at 4C.
7. Remove primary antibody and replace with PBS. Wash three times for 10 minutes each.
8. Add secondary antibody at appropriate dilution. Incubate for 1 hour at room temperature.
9. Remove secondary antibody and replace with PBS. Wash three times for 10 minutes each.
10. Counterstain DNA with DAPI if required.





### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NLS1878**

---

NLS1878PEP	FPRL1/FPR2 Antibody Blocking Peptide
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NLS1878](http://www.novusbio.com/reviews/submit/NLS1878)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

